

For audio, please call:  
866-737-4154  
Passcode: 6437042#



California Landscape Conservation Cooperative  
Webinar Series

CaliforniaLCC.org

Today's guest presenter – Dr. Kristin Byrd, US Geological Survey

Topic: “Climate Change and Land Use Scenarios for Habitat Threat Assessments on California Rangelands”

Webinar will begin shortly. Please mute your phone.

March 14, 2013

# Climate change/land use change scenarios for assessing threats to ecosystem services on California rangelands

Kristin Byrd<sup>1</sup>, Lorraine Flint<sup>2</sup>, Frank Casey<sup>3</sup>, and Pelayo Alvarez<sup>4</sup>



March 14, 2013

<sup>1</sup>USGS Western Geographic Science Center, Menlo Park, CA,

<sup>2</sup>USGS California Water Science Center, Sacramento, CA ,

<sup>3</sup>USGS Science and Decisions Center, Reston, VA,

<sup>4</sup>Defenders of Wildlife, Sacramento, CA,


# Ecosystem services provided by rangelands

- Food, fiber and fuel
- **Wildlife habitat**
- **Water**
- **Carbon sequestration**
- Adaptation to climate change
- Open space, cultural values



# Integrated Threats to Rangelands

- In California 20,000 acres of rangelands are lost every year
- Privately owned
- Cattle ranching: low profits
- Low levels of protection



Land conversion and climate change lead to loss of grazing land, water availability, and altered species distribution

# Rangeland Coalition Focus Area Map (TNC, 2007)

<http://www.carangeland.org/focusarea.html>

Dark blue: Critical Conservation Areas

(Privately-owned rangelands that have high biodiversity value and require conservation action in the next 2-10 years.)

## CALIFORNIA RANGELAND CONSERVATION COALITION



### Focus Area Prioritization

Rangelands represent one of the most threatened habitats throughout the western United States. In addition to being threatened, these oak savannah and grassland habitats have relatively low levels of conservation management while maintaining high biodiversity values. Many grassland birds, native plants, and threatened vernal pool species on the landscape benefit from responsible grazing practices. Instead, privately-owned rangelands face threats from increased low density, rural residential housing development in the foothills and conversion to other uses.

Out of this concern environmentalists, cattlemen and government agencies have come together to form a sustainability conservation partnership, the California Rangeland Conservation Coalition. This map illustrates the coalition's priority focus areas for conservation and enhancement. The Rangeland Coalition works with willing private landowners to preserve rangelands through conservation easements and to carry out habitat enhancement projects for common and threatened species.

Conservation Easements include owner:  
Private Land  
Conservation Easement  
Conservation Easement  
Conservation Easement

Scale: 0 100 200 Miles



**Priority Areas**

- Area Critical to CRC's Conservation Goals
- Area Important to CRC's Conservation Goals
- Important or Critical Area Threatened by Regeneration\*

**Non-Priority Areas**

- Currently (2004) Urban or Suburban
- Area Facing Regeneration Threat outside of Priority Areas by 2020

**Other Features**

- CRC's Boundary
- Additional Privately Protected Land\*\*
- Lake and Reservoirs

\* Areas facing regeneration are those areas that are currently urban or suburban and are projected to be converted to rangeland by 2020. \*\* Areas that are currently privately protected but not included in the CRC's focus area.





# Project Goals

- Six spatially-explicit climate change/land use change scenarios from years 2000 – 2100 consistent with three IPCC emission scenarios and two global climate models –

## B1 (sustainability)

1. PCM (warm, wet future)
2. GFDL CM 2.1 (hot, dry future)

## A1B (wealth and technology)

1. CSIRO Mark 3.5 GCM (warm, wet future)
2. MIROC 3.2 (medres) (hot, dry future)

## A2 (population pressures)


1. PCM (warm, wet future)
2. GFDL CM 2.1 (hot, dry future)

- Assess potential threats to rangeland ecosystem services
  1. wildlife habitat
  2. water availability (Lorraine Flint and Alan Flint, USGS)
  3. carbon sequestration



# Project Goals, continued

3. An economic analysis of scenarios to quantify economic costs and benefits (Frank Casey, USGS)
4. A web-based visualization tool, and
5. An outreach program that will target the Rangeland Coalition network to communicate how results can be applied to conservation and land management decisions. (Pelayo Alvarez, Defenders of Wildlife)




# Why IPCC emission scenarios?

- Climate scenarios and land use scenarios need to be **logically consistent** to form the basis for integrated assessments and long-term policies (Bierwagen et al. 2010).
- Existing land-use land-cover (LULC) change modeling and downscaled global climate models based on the same scenarios – **A1B, A2, B1**
  - USGS LULC change scenarios
  - USGS ensemble projections of climate and hydrology for California (Lorraine Flint and Alan Flint, USGS)



















# National Assessment of Ecosystem Carbon Sequestration and Greenhouse Gas Fluxes

[http://www.usgs.gov/climate\\_landuse/land\\_carbon/](http://www.usgs.gov/climate_landuse/land_carbon/)



## USGS National Land Cover Dataset (NLCD)

### Land use/land cover classes class name

|   |                    |
|---|--------------------|
|    | Agriculture        |
|    | Barren             |
|    | Deciduous Forest   |
|    | Developed          |
|    | Evergreen Forest   |
|    | Grassland          |
|    | Hay/Pasture        |
|    | Herbaceous Wetland |
|   | Mech Disturbed NF  |
|  | Mech Disturbed OP  |
|  | Mech Disturbed PVT |
|  | Mining             |
|  | Mixed Forest       |
|  | Shrubland          |
|  | Water              |
|  | Woody Wetland      |

- Three LULC change scenarios for each EPA Level III ecoregion (Ben Sleeter, USGS)
- FORE-SCE model: maps of LULC change by scenario/year (Terry Sohl et al., USGS)
- GEMS biogeochemical model: annual total ecosystem carbon change per LULC class (S. Liu et al., USGS)

# Driving Force Assumptions for the United States based on IPCC Emission Scenarios

(table adapted from Ben Sleeter, USGS)

|                                 | <b>A1B – wealth and technology</b>                                | <b>A2 – population pressures</b>                                 | <b>B1 - sustainability</b>                                       |
|---------------------------------|---|--|--|
| <b>DEMOGRAPHICS</b>             | Medium growth, sprawl   | High growth, sprawl  | Medium growth, densification                                     |
| <b>ECONOMICS</b>                | Very High Income  | Medium Income  | High Income  |
| <b>TECHNOLOGY</b>               | Very High rate of innovation                                      | Low rate of innovation   | High rate of innovation  |
| <b>ENERGY</b>                   | Balanced between several sources                                  | Fossil fuel intensive  | Rapid diffusion of “green” energy resources                      |
| <b>CLIMATE</b>                  | Temperature change, best estimate and range: 2.8 °C; 1.7 – 4.4 °C | Temperature change, best estimate and range: 3.4 °C; 2.0 – 5.4°C | Temperature change, best estimate and range: 1.8 °C; 1.1 – 2.9°C |
| <b>ENVIRONMENTAL PROTECTION</b> | Mixed-use based conservation                                      | Conservation lower priority                                      | Conservation high priority                                       |

# Scenario Narratives for CA Rangelands




Rancher's Focus Group, January 2012, Davis CA

## Key Concerns about ranching future:

- Limited availability of grazing land for lease
- Fragmentation of grazing land
- Forage quality and quantity
- High start-up investment

# Scenario Narratives for CA Rangelands

## – Alternative conservation plans




# Integrated Scenarios

Three IPCC scenarios  
(A1B, A2, B1)  
Two climate models  
(warm, wet or hot, dry)


Land use/land cover  
change +

Climate/hydrology  
decadal change

Maps by scenario/year  
to 2100 at ~250 meter  
resolution




California Rangeland  
Conservation Coalition  
Focus Area




EPA Level III Eco-regions:  
Central Valley and  
Chaparral and Oak Woodlands


# Land-use land-cover change 2006 to 2100; B1, A2, A1B




Present-Day






PCM B1 2100  
GFDL B1 2100



PCM A2 2100  
GFDL A2 2100



CSIRO A1B 2100  
MIROC A1B 2100



Land use-land cover/  
Climate/Hydrological  
Change

Precipitation  
Minimum Winter Temp.  
Maximum Summer Temp.  
Climatic Water Deficit  
Potential  
Evapotranspiration  
Decadal averages 2010 –  
2100, 250 meters


FORE-SCE LULC  
Change Model  
Annual maps of land use  
change 2006-2100, 250  
meters

Ecosystem Services  
Change (water, carbon,  
habitat)

Basin Characterization  
Model  
Runoff, Recharge, Stream  
Discharge  
2010, 2040, 2070, 2100

Change to Priority  
Conservation Areas (TNC,  
2007)  
Decadal change 2010 –  
2100

GEMS biogeochemical  
model  
Total Ecosystem Carbon  
2006 – 2050




## Landscape-level analysis

- Land use/climate change for conservation scenarios
- Water-wildlife hotspots




# LULC change in B1 conservation areas


## A2 Scenario




## B1 Scenario




# Summer maximum temperature by scenario, B1 conservation areas






## LULC change in A1B conservation areas





## Summer maximum temperature by scenario, A1B conservation areas




# Water-wildlife hotspots for dry scenarios (draft)



Percent change in water availability (runoff+recharge) by basin from 1951-1980 to 2071-2100



Percent change in water availability in basins with >25% loss in critical habitat



## Case Study of Six Watersheds:

### North:

Upper Stony  
Lower Butte

### Central:

Lower Cosumnes  
Alameda Creek

### South:

Upper Tule  
Estrella


Changes in:

- Wildlife habitat
- Carbon
- Runoff, recharge, streamflow




# Change in Total Ecosystem Carbon - Upper Stony Watershed

## Grassland




## Combined Forest




# Habitat Change - Upper Stony Watershed


## A1B



## A2




## B1




- Developed
- Agriculture
- Hay/Pasture
- Grassland
- Herbaceous Wetland
- Woody Wetland
- Shrubland
- Forest

# Change in Total Ecosystem Carbon – Consumnes Mokelumne Watershed

## Grassland




## Combined Forest




# Habitat Change – Consumnes Mokelumne Watershed


## A1B



## A2




## B1




- Developed
- Agriculture
- Hay/Pasture
- Grassland
- Herbaceous Wetland
- Woody Wetland
- Shrubland
- Forest

# Change in Total Ecosystem Carbon – Alameda Creek Watershed

## Grassland




## Combined Forest




# Habitat Change – Alameda Creek Watershed


## A1B




## A2



## B1




- Developed
- Agriculture
- Hay/Pasture
- Grassland
- Herbaceous Wetland
- Woody Wetland
- Shrubland
- Forest






# Upper Stony Watershed

## Recharge/Runoff




## Streamflow




# Alameda Creek Watershed

## Recharge/Runoff




## Streamflow




# Consumnes Mokelumne Watershed

## Recharge/Runoff



## Streamflow





# Summary

- Potential for C sequestration decreases with area and rate of grassland conversion
- The ratio of recharge to runoff decreases with increasing urbanization (Alameda, Cosumnes)
- Amount of change depends on current soil storage capacity, more change if urbanization on deep soils
- In non-urbanized watersheds, ratio of recharge to runoff can increase in dry years (Upper Stony)
- Has implications on water resource planning – water supply and habitat and need to plan for extreme events



# Social value of carbon : avoided marginal damages from carbon emissions to a society as a whole

- Value: \$45/ton CO<sub>2</sub>e (Kroeger 2012)
- Carbon: 47 t CO<sub>2</sub>e/acre (Koteen et al. 2005, Silver et al. 2010)

| Watershed           | Grassland acreage lost from 2010 to 2050 | Social value of carbon stocks (CO <sub>2</sub> e) |
|---------------------|--|---|
| Upper Stony         |  |   |
| A1B                 | 14429                                    | \$30,517,335.00                                   |
| A2                  | 8417                                     | \$17,802,012.12                                   |
| B1                  | 12120                                    | \$25,633,800.00                                   |
| Alameda Creek       |  |   |
| A1B                 | 66749                                    | \$141,174,135.00                                  |
| A2                  | 45699                                    | \$96,653,385.00                                   |
| B1                  | 48062                                    | \$101,651,130.00                                  |
| Consumnes-Mokelumne |  |   |
| A1B                 | 62765                                    | \$132,747,975.00                                  |
| A2                  | 23768                                    | \$50,269,320.00                                   |
| B1                  | 13328                                    | \$28,188,720.00                                   |

# Future Economic Analysis

- Link decreases in recharge to costs associated with less availability of water for consumptive and environmental uses
- Link increases in run-off to potential costs associated with mitigating increased sedimentation and other water quality issues
- Analyze changes in stream flow with respect to economic impacts on aquatic habitat
- Analyze land use changes with respect to potential economic impacts on wildlife habitats, including use and non-use values

# Outreach

## a) Key messages:


- Inform stakeholders of impacts of climate change and land use change to rangeland ecosystem services
- Decision-making tool for prioritization of climate change mitigation strategies (i.e restoration sites, conservation easements)
- Raise awareness about the importance of rangelands in providing ecosystem services

## b) Targets

- Ranches and land managers
- Government agencies
- Non-profits: Ag and conservation organizations
- Others: researchers, planners, legislators, general public



# Acknowledgments



Chris Souldard, Western Geographic Science Center, Menlo Park, CA

Adam McClure, Western Geographic Science Center, Menlo Park, CA

Zhiliang Zhu, LandCarbon Project Lead, Reston, VA

Shuguang Liu, EROS Data Center, Sioux Falls, SD

Terry Sohl, EROS Data Center, Sioux Falls, SD

Ben Sleeter, Western Geographic Science Center, Menlo Park, CA

The USGS LandCarbon Team

Funding by the California Landscape Conservation Cooperative





California Landscape Conservation Cooperative  
Webinar Series

CaliforniaLCC.org

**Thank you for joining us. A recorded version of this webinar will be available on our website in about a week.**

If you have questions about the webinar, contact Rebecca Fris at 916-278-9415.

March 14, 2013